☐ POV Age ____ Sex ○ Male ○ Female Activation Time ____

Level One:

Physiological Criteria

Pediatric (Age 13 Years and Below)

☐ Capillary Refill: >2 Seconds with Signs of Trauma

☐ AVPU: Responsive to Pain or Unresponsive

Age	SBP or RR* Indicators	
< 1 Year	☐ SBP <60mmHg	☐ RR* <10 or >50
1-10 Years	☐ SBP <(70+2*Age)	☐ RR* <10 or >40
>10 Years	☐ SBP <90mmHg	☐ RR* <10 or >30
signs of respiratory insufficiency such as hypoxia, grunting, or accessory muscle use		

Adult (Age of 14 Years or Older)

☐ SBP <90mmHg

☐ RR <10 or >29

☐ GCS <=13 with known trauma mechanism

Geriatric (Age 65 or older)

☐ Two or More "Level Two" Criteria (Mark the additional criteria)

Anatomical Criteria

☐ Airway instability/compromise or prehospital intubation

☐ Transferred trauma patients receiving blood products to maintain vital signs

☐ Paralysis or Suspected C/T/L-Spine Injury with Neurologic Deficit

☐ Pulseless Extremity Due to Trauma

☐ Two or More Proximal Long Bone Fractures

☐ Penetrating Injury to Head, Neck, Chest, Groin, Abdomen or Proximal to the Knee or Elbow

☐ Traumatic Extremity Amputations Above Knee or Elbow

☐ Open and/or Depressed Skull Fracture

☐ Unstable Pelvic Fracture

☐ Severe Chest Injury and/or Flail Chest

☐ Positive FAST exam

☐ **Trauma Surgeon or ED Physician Request**

Name: ____

Level Two:

Physiological Criteria

☐ GCS=14 with known trauma mechanism

Mechanism of Injury

Auto

☐ Pedestrian/bicyclist hit by auto >20 MPH, run over or thrown >15ft.

☐ Death of an occupant in same vehicle

☐ Ejection or Separation from auto, motorcycle, horse, ATV etc.

☐ Motorcycle crash >20 MPH

☐ Major Auto Accident Suggested by Prolonged Extrication (>20min),

☐ Intrusion into the passenger compartment >12 inches

☐ Signs of abdominal injury (seatbelt sign, distension, firmness)

Falls

☐ Adults >15 ft (one story=10ft)

☐ Pediatrics >10 ft or 2 times the height of the child

☐ Fall from any height on blood thinners with evidence of Head Trauma

Burn

☐ 2nd and 3rd degree burns > 10% BSA

☐ Inhalation injury

☐ Electrical burn

Other Considerations:

☐ Suspected Spinal Injury Without Neurologic Deficit

☐ Pregnancy > 20 Weeks

☐ Suspected Non-Accidental Trauma (e.g. Child Abuse)

☐ Drowning/Asphyxiation

☐ EMS Request

☐ **Trauma Surgeon or ED Physician Request**

Name: ____

Patient Label